



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

with the condition

$$\lambda_1 + \mu_1 + \nu_1 = 0, \quad (3)$$

and that of a second plane embracing (1) is

$$\frac{(x-a)\lambda_2}{l} + \frac{(y-b)\mu_2}{m} + \frac{(z-c)\nu_2}{n} = 0, \quad (4)$$

with the condition

$$\lambda_2 + \mu_2 + \nu_2 = 0. \quad (5)$$

If (2) and (4) are orthogonal,

$$\frac{\lambda_1}{l} \cdot \frac{\lambda_2}{l} + \frac{\mu_1}{m} \cdot \frac{\mu_2}{m} + \frac{\nu_1}{n} \cdot \frac{\nu_2}{n} = 0, \quad (6)$$

and if (2) and (4) touch

$$a_1x^2 + b_1y^2 + c_1z^2 = 1, \quad (7)$$

$$\left(\frac{\lambda_1}{l}\right)^2/a_1 + \left(\frac{\mu_1}{m}\right)^2/b_1 + \left(\frac{\nu_1}{n}\right)^2/c_1 = \left(\frac{a\lambda_1}{l} + \frac{b\mu_1}{m} + \frac{c\nu_1}{n}\right)^2, \quad (8)$$

and

$$\left(\frac{\lambda_2}{l}\right)^2/a_1 + \left(\frac{\mu_2}{m}\right)^2/b_1 + \left(\frac{\nu_2}{n}\right)^2/c_1 = \left(\frac{a\lambda_2}{l} + \frac{b\mu_2}{m} + \frac{c\nu_2}{n}\right)^2. \quad (9)$$

Now (3), (5), (6), (8), (9) is a system of *five* equations for the determination of the *six* ratios  $\lambda_1/\nu_1, \mu_1/\nu_1; \lambda_2/\nu_2, \mu_2/\nu_2$ ; and  $l/n, m/n$ , giving an indeterminate solution. The values of  $l/n, m/n$  are the only ones needed in (1).

There seems to be a missing condition in the statement of the problem.

If the direction of the line (1) were constant, or  $l : m : n$ , constant, the *locus* of the line (1) would be a right circular cylinder, or, in other words, the locus of the line of intersection of constant direction of pairs of orthogonal tangent planes to a central conicoid is a right circular cylinder.

*Note.*—The single condition on the line restricts it to be a member of a line complex whose order apparently may be as high as 8. It would be desirable to determine this explicitly. EDITORS.

#### 497. Proposed by NATHAN ALTSHILLER, University of Oklahoma.

Find the locus of the mid-point of the segment determined by two spheres on any line passing through any point common to the two spheres.

#### SOLUTION BY S. W. REAVES, University of Oklahoma.

Let the plane of the common circle of the spheres be chosen for  $yz$ -plane, and the line of centers for  $x$ -axis. Let the radius of the common circle be  $k$ , and let  $(a, 0, 0)$  and  $(b, 0, 0)$  be the centers of the two spheres. Then the equation of one sphere is

$$(x-a)^2 + y^2 + z^2 = a^2 + k^2, \quad (1)$$

or

$$x^2 + y^2 + z^2 - 2ax = k^2;$$

and, similarly, the equation of the other is

$$x^2 + y^2 + z^2 - 2bx = k^2. \quad (2)$$

Let  $l, m, n$  be the direction cosines of an arbitrary line through the point  $M(0, 0, k)$ . Then the equation of the line may be written

$$\frac{x}{l} = \frac{y}{m} = \frac{z-k}{n} = r, \quad (3)$$

where  $r$  is the length of the segment joining  $(0, 0, k)$  and  $(x, y, z)$ .

To find the length of the segment  $MP$  cut from this line by the sphere (1), we substitute  $x = lr, y = mr, z = nr + k$  in equation (1) and solve for that value of  $r$  which is not zero. We thus find at once

$$r = MP = 2al - 2kn.$$

Substituting the same values in (2), we find likewise for the segment  $MQ$  intercepted by the

second sphere,

$$MQ = 2bl - 2kn.$$

If  $R$  be the mid-point of  $PQ$ , then

$$MR = \frac{MP + MQ}{2} = (a + b)l - 2kn.$$

Hence, if in (3) we set

$$r = (a + b)l - 2kn, \quad (4)$$

we readily obtain expressions for the coördinates  $x, y, z$  of the middle point  $R$  of the segment determined by the two spheres on the line through  $(0, 0, k)$  and having the direction  $(l, m, n)$ . To find the equation of the locus of  $R$  for all directions of the line we must eliminate  $l, m, n$ , which we do as follows:

From (3), we have,

$$x^2 + y^2 + (z - k)^2 = r^2; \quad (5)$$

and from (4) and (3),

$$r = (a + b) \left( \frac{x}{r} \right) - 2k \left( \frac{z - k}{r} \right),$$

or

$$r^2 = (a + b)x - 2k(z - k). \quad (6)$$

Equating the values of  $r^2$  given by (5) and (6), we have for the equation of the locus of  $R$ ,

$$x^2 + y^2 + z^2 - (a + b)x = k^2. \quad (7)$$

The locus is, therefore, a sphere whose center is midway between the centers of the given spheres and which contains all points common to these spheres. It is evident then that the choice of any point other than  $M$  common to the two spheres would lead to the same locus.

Also solved by the PROPOSER.

### CALCULUS.

#### 413. Proposed by OSCAR S. ADAMS, Washington, D. C.

Determine a function of  $x$  independent of  $b$ , such that

$$\int_b^{b+1} f(x)dx = \frac{1}{b+1},$$

the real part of  $b$  being positive.

SOLUTION BY C. F. GUMMER, Kingston, Ontario.

There is one solution only satisfying the conditions:

- (i)  $f(x)$  is one-valued and continuous in the right half plane of complex numbers, and
- (ii)  $f(x + n)$  approaches 0 as  $n$  approaches  $+\infty$  through integral values for every  $x$ .

Let  $f(x)$  be such a solution. Differentiating the equation

$$\int_b^{b+1} f(x)dx = \frac{1}{b+1}, \quad (1)$$

we get, after substituting  $x$  for  $b$ ,

$$f(x+1) - f(x) = -\frac{1}{(x+1)^2}.$$

Similarly,

$$\begin{array}{rcl} f(x+2) - f(x+1) & = & -\frac{1}{(x+2)^2}, \\ \vdots & & \vdots \\ f(x+n) - f(x+n-1) & = & -\frac{1}{(x+n)^2}. \end{array}$$

Adding these equations, we have